

Empowering Your Recovery After Chemotherapy

Naturally Rejuvenating, Restoring, and Rebuilding to Prevent Recurrence.





Product Name

Purobest

Berberine LPX 250

250 ma of Std to 97% Berberine

Purobest

Green Tea SPX 500

500 mg of EGCG from Green Tea (Camellia Sinensis) Extract std to 40%

Purobest

Curcu LPX 2500

2500 mg of 95% Curcuminoids



Main Action

Berberine exhibits Anti-cancer effects across various cancer types by inhibiting proliferation, suppressing metastasis, and inducing apoptosis through interactions with microRNAs and inhibition of the BCL-2/BAX signaling pathway. Potentially involves overcoming multidrug resistance (MDR) and improving the efficacy and safety of existing cancer treatments.

EGCG induces apoptosis and cell growth arrest by inhibiting MAPK, activating FOXO genes, and reducing ET-1 expression through **Akt- and AMPK-stimulated** FOXO1 regulation, involving key proteins like calmodulin and multiple pathways such as angiogenesis and AP-1 inhibition. EGCG causes modulation of genes, including FOXO, and inhibition of proteins like AP-1, impacting pathways crucial for cell survival and angiogenesis.

Curcumin upregulates E-cadherin (adhesion promoter) and downregulates vimentin & N-cadherin (migration enhancers), restoring cell connections and hindering escape. It triggers programmed cell death in chemotherapy resistant cells, eliminating them and reducing overall tumor burden.



Specific benefits

It benefits cancer treatment, inhibits metastasis, induces apoptosis, and overcomes resistance.

It induces apoptosis, arrests cell growth, and modulates key genes and pathways, potentially enhancing treatment outcomes by targeting multiple aspects of cancer.

Stimulates cell death in chemotherapy-resistant cancer cells.

Not recommended for



When to **Prescribe**

For recurrence avoidance post treatment and for improved efficacy during treatment

Advised for cancer survivors after chemotherapy for recurrence avoidance.

Recommended for cancer survivors after chemotherapy for recurrence avoidance

individuals with severe bleeding

disorders or severe gastric

before and two days after

chemotherapy.

15-30 minutes

issues. Avoid using one day



Major

Not recommended for Contraindications individuals with severe bleeding disorders or severe gastric issues. Avoid use one day before and two days after chemotherapy.

Not recommended for individuals with severe bleeding disorders or severe gastric issues. Avoid use one day before and two days after chemotherapy.

> 1 Sachet OD diluted in 300 ml of water and taken over



Dosage

1 Capsule BD

1 Tablet OD







